GENITOURINARY MEDICINE

Editor: A Mindel (Sydney)

President MSSVD: JRW Harris (London) Hon Sec MSSVD: T McManus

G Bird (Edinburgh)
D A Cooper (Sydney)
L Corey (Seattle)
R A Coutinho (Amsterdam)
B A Evans (London)
J S Greenspan
(San Francisco)

D J Jeffries (London)
F Judson (Denver)
C J N Lacey (Leeds)
J M A Lange (Amsterdam)
A G Lawrence (London)
A Méheus (Geneva)
A R Moss (San Francisco)

Assistant Editor: C A Carne (Cambridge)
Assistant Editor, Abstracts: C Sonnex (Cambridge)

R S Pattman
(Newcastle-upon-Tyne)
P Piot (Antwerp)
M Stanley (Cambridge)
D Taylor-Robinson
(Harrow)
J Weber (London)

I V D Weller (London)
L Weström (Lund)
H Young (Edinburgh)
A J Zuckerman (London)
H zur Hausen (Heidelberg)
Editor British Medical Journal
Technical Editor: Kenneth Till
(Taunton, UK) Fax 09847-315

This Journal, founded by the Medical Society for the Study of the Venereal Diseases, publishes original work on the investigation and treatment of genitourinary and allied disorders, and review articles, correspondence, and abstracts.

ADVICE TO AUTHORS Papers for publication, which will be accepted on the understanding that they have not been and will not be published elsewhere and are subject to editorial revision, should be sent in duplicate to Prof A Mindel, Academic Department of Sexual Health Medicine, Sydney Hospital, Macquarie St, Sydney 2000, GPO Box 1614, Sydney, NSW, Australia. Tel: 61 2228 2111; Fax: 61 2222 3186. All authors must give signed consent to publication. Submitted papers must be accompanied by the following statement, signed by all the authors: "No paper resembling the enclosed article has been or will be published except in Genitourinary Medicine. We transfer all copyright ownership to Genitourinary Medicine." Manuscripts will only be acknowledged if a stamped addressed postcard or international reply coupon is enclosed. Rejected papers will not be returned. Figures will be returned if requested at the time of submission.

Full details of requirements for manuscripts in the Vancouver style (BMJ 1982;284:1766-70) are given in Uniform requirements for manuscripts submitted to biomedical journals, available from the Publishing Manager, British Medical Journal, BMJ Publishing Group, BMA House (50p post free). Briefly details are as follows:

- (1) SCRIPTS (including correspondence and book reviews) must be typewritten on one side of the paper in double spacing with ample margins. Two copies should be sent.
- (2) EACH SCRIPT should include, in the following order: a brief summary (structured summaries are preferred), typed on a separate sheet, outlining the main observations and conclusions; the text divided into appropriate sections; acknowledgements; references; tables, each on a separate sheet; and legends for illustrations.
- (3) THE TITLE of the paper should be as brief as possible.
- (4) THE NUMBER OF AUTHORS should be kept to the minimum, and only their initials and family names used.
- (5) ONLY THE INSTITUTION(s) where work was done by each author should be stated.
- (6) SI UNITS must be used. If old fashioned units are used, SI units should be given in parentheses or, for tables and figures, a conversion factor given as a footnote.
- (7) ONLY RECOGNISED ABBREVIATIONS should be used.
- (8) ACKNOWLEDGEMENTS should be limited to workers whose courtesy or help extended beyond their paid work, and supporting organisations.
- (9) FIGURES should be numbered in the order in which they are first mentioned in the text. Captions should be typed on a separate sheet. DIAGRAMS: use thick, white paper and insert lettering lightly in pencil. PHOTOGRAPHS: should be marked lightly on the back with the author's name and indicating the top, and should not be attached by paper clips or pins. They should be trimmed to include only the relevant section (and will be reproduced 68 or 145 mm wide) to eliminate the need for reduction. Photomicrographs must have internal scale markers. Radiographs should be submitted as photographic prints, carefully prepared so that they bring out the exact point to be illustrated.
- (10) TABLES should be numbered, have titles, and be typed on separate sheets. Please avoid large tables and use the format which appears in current issues.

(11) REFERENCES should be numbered consecutively the first time they are cited and identified by arabic numbers in the text, tables, and legends to figures. Authors must take full responsibility for the accuracy of their references, and the list should be kept as short as practicable. It should be in the order in which references are first mentioned, and should include (in the following order), journals: author's name and initials, title of paper, name of journal (in full or abbreviated according to the list in *Index Medicus*), year of publication, volume number, and first and last page numbers; books: author's name and initials, full title, edition, place of publication, publisher, and year of publication. When a chapter in a book is referred to, the name and initials of the author of the chapter, title of the chapter, "In:", name and initials of the editor, and "ed" should precede book title, etc as above. In references to journals or books, when there are seven or more authors the names of the first three should be given followed by "et al." Names of journals no longer published or not in *Index Medicus* should be given in full - for example, British Journal of Venereal Diseases.

PROOFS Contributors receive one proof, and should read it carefully for printers' errors and check the tables, figures, legends, and any numerical, mathematical, or other scientific expressions. Alterations should be kept to a minimum.

REPRINTS A limited number of reprints may be ordered at cost from the Publishing Manager when the proofs are returned.

NOTICE TO SUBSCRIBERS This Journal is published bimonthly. The annual subscription rates are £119.00 and \$216 in the USA. Subscribers may pay for their subscriptions by Access, Visa, or American Express by quoting on their orders the credit or charge card preferred, appropriate personal account number, and the expiry date of the card. Orders should be sent to the Subscriptions Manager, Genitourinary Medicine, BMJ Publishing Group, BMA House, Tavistock Square, London WC1H 9JR. Orders can also be placed locally through any leading subscription agent or bookseller. (For the convenience of readers in the USA, subscription orders, with or without payment, can be sent to: British Medical Journal, Box 560B, Kennebunkport, Maine 04046. All inquiries, however, including those regarding air mail rates and single copies already published, should be addressed to the publisher in London.)

Second class postage paid, Rahway NJ Postmaster. Send address changes to: *Genitourinary Medicine* c/o Mercury Airfreight International Ltd Inc, 2323 Randolph Avenue, Avenel, NJ 07001, USA.

NOTICE TO ADVERTISERS Applications for advertisement space and for rates should be addressed to the Advertisement Manager, *Genitourinary Medicine*, BMJ Publishing Group, BMA House, Tavistock Square, London WC1H 9IR.

COPYRIGHT © 1993 by Genitourinary Medicine. This publication is copyright under the Berne Convention and the International Copyright Convention. All rights reserved. Apart from any relaxations permitted under national copyright laws, no part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the prior permission of the copyright owners. Permission is not, however, required to copy abstracts of papers or of articles on condition that a full reference to the source is shown. Multiple copying of the contents of the publication without permission is always illegal.

ISSN 0266-4348

Published by BMJ
Publishing Group,
and printed by
Eyre & Spottiswoode Ltd,
London and Margate

158 Matters arising

MATTERS ARISING

Actinomyces israelii in the female genital tract

I have been reading with interest Dr Evans' review of Actinomyces israelii in the female genital tract. I am troubled by his suggestion that the IUCD should be removed in symptomless patients. Sperm remains viable in the female genital tract for at least 5 days and I would recommend that if intercourse has taken place in the past week, another method of contraception should be provided and the IUCD removed when there is no longer a risk of pregnancy or that the IUCD be removed immediately and replaced by another.

In the few cases I have seen and treated by the above methods, cervical smears taken at 3 months have all been actinomycosis free.

ELIZABETH MAYALL Margaret Jackson Centre, 4 Barnfield Hill, Exeter, EX5 4AA, UK

1 Evans DTP. Actinomyces in the female genital tract. Gen Med 1993;69:54-9.

The importance of ethnicity as a risk factor for STDs and sexual behaviour

A number of papers have addressed the relevance of race, country of birth and "ethnicity" in the epidemiology of sexually transmitted diseases. A recent example is the paper by Hooykaas et al.1 Here the authors speak of Turkish men, Surinam men, Dutch men. They say that the country of birth was a strong predictor of STDs. We feel that the country of birth, nationality and ethnicity are not surrogates for one another. Sexual mores are often culturally modulated and religion, and upbringing are factors worth studying. We suggest that: firstly, the terminology employed should be defined by the researchers if it is not in accord with a standard dictionary (eg, the Shorter Oxford) and secondly, religion (even if only nominal) should be recorded when collecting the data. Some Surinam people, for example, are Creoles, others are Hindustanees and so on. Their religions include Christianity, Islam, Hinduism. Men born in Turkey would fall in to two main groups: Turks and Kurds. The Kurds may well exhibit a different pattern of sexual mores from the Turks. We believe that a study of the cultural and religious characteristics of the patients recruited in similar studies would be more rewarding than the analyses published so far.

J K ANAND 68 Ledbury Rd, Peterborough PE3 9PJ

A DE LA COURT St Mary's Hospital, London W2 1PG, UK.

1 Hooykaas C, van der Velde FW, van der Linden MMD, van Doornum GJJ, Coutinhou RA. The importance of ethnicity as a risk factor for STDs and sexual behaviour among heterosexuals. Genitourin Med 1991;67:378-83.

NOTICES

11th Regional Conference of Dermatology

This will be held 21-24 May 1994 in Singapore. The theme will be *Trends and Recent Developments in Dermatology*. Information may be obtained from: The Secretariat, 11th Regional Conference of Dermatology, c/o Conference and Exhibition Management Services Pte Ltd, 09-43 World Trade Centre, Republic of Singapore 0409, Attention Maggie Phang.

The Medical Society for the Study of Venereal Diseases (MSSVD) Undergraduate Prize Regulations

- 1 A prize of £200.00 to be called the MSSVD Undergraduate Prize, will be awarded annually by the MSSVD (provided an entry of a suitable standard is received).
- 2 Entries for the prize will take the form of a report written in English.
- 3 The subject of the report should be related to sexually transmitted disease, genitourinary medicine or HIV infection.
- 4 The report should concern original and unpublished observations made by the entrant. The report, which should not exceed 2,000 words, should include an introduction to the subject, methods used to make the observations, findings and discussion. A summary of the report on a separate sheet should also be provided. Entries must be machine or type-written and double spaced on one side only of A4 paper. Three copies must be submitted.
- 5 The subject must be approved by a genito-urinary physician to the entrant's medical school. The observation must be made before full registration. A winner may not enter for the Prize again. Each entry should be accompanied by a declaration that these conditions have been fulfilled.
- 6 Entries should be submitted to the Hon Secretary of the Medical Society for the Study of Venereal Diseases by 30th June each year. They will then be considered by the President, the Hon Secretary and the Hon Treasurer. When appropriate other experts may be consulted. These assessors will make recommendations to Council who will make the final decision concerning the Prize.
- 7 Entries must be submitted within 12 months of full registration or its equiva-
- 8 Regulations are obtainable from the Hon Secretary MSSVD.
- 9 The assessors may ask the editor of an appropriate journal to consider an entry for publication. If so, it will be received for publication in the usual way.

BOOK REVIEW

Colour guide—HIV infection and AIDS. By MA Birchall, SM Murphy, Edinburgh, Churchill Livingstone. (£7.95: pp 121). 1992. ISBN 0-4430-4578-X.

The authors of this slim paper back have succeeded in providing an easy to read book containing a lot of useful information. For its size, the book is surprisingly comprehensive, covering a wide spectrum of diseases related to HIV infection. The diagnosis, clinical features and management of each condition is discussed with sections on pathology and aetiology where appropriate. The majority of the slides are excellent with some old favourites but a number of new slides.

I am unclear to whom the authors have pitched this book. If as stated on the back cover the market is "students and health care specialists in all fields of medicine" then it would have been helpful to include a list of abbreviations at the front to avoid hunting through the text to find the meaning of BAL, GIT, MAI, CMV etc. There is also no indication in most chapters of what are the common or rare presentations of HIV disease. Six of 26 chapters are dedicated to otolaryngology with three separate chapters, albeit short, on the ear (would this have any connection with the specialist area of one of the authors?!). Arrows indicating the abnormalities mentioned in the captions of the histology and electron micrography slides would be very helpful for non-special-

The authors devised a logical list of contents. However, the chapter on "Early clinical features" left me confused. It is not clear whether the chapter refers to early clinical features of AIDS or HIV infection. I presumed the latter but references made to the onset of late disease and the inclusion of a section on oral candidiasis in this chapter left me in doubt. The index was not comprehensive. In particular some therapeutic agents mentioned in the text were omitted.

These are just minor criticisms of an otherwise well presented and useful guide. As a specialist in the field, I would see no reason to buy it but as an introduction to the topic, for quick revision and painless learning it certainly fits the pocket.

A ROBINSON

CURRENT PUBLICATIONS

Selected titles from recent reports published worldwide are arranged in the following sections:

Gonorrhoea Chlamvdia Pelvic inflammatory disease Candidiasis Bacterial vaginosis Trichomoniasis Syphilis and other treponematoses Herpes Human papillomavirus infection Cervical cytology and colposcopy Other sexually transmitted diseases Public health and social aspects Microbiology and Immunology Dermatology Miscellaneous

Gonorrhoea

Fulminant endocarditis due to infection with penicillinase-producing Neisseria gonorrhoeae

PJ WEISS, CA KENNEDY, DF MCCANN, HE HILL, EC OLDFIELD, Sex Transm Dis 1992;19:288.

Evaluation of new anti-infective drugs for the treatment of uncomplicated gonorrhoea in adults and adolescents

HH HANDSFIELD, JA MCCUTCHAN, L COREY, AR RONALD, Clin Infect Dis 1992;15:S123.

Evaluation of treatment with single-dose ampicillin/sulbactam with probenecid or ceftriaxone in patients with uncomplicated gonorrhoea

LM BADDOUR, L BUSBY, E SHAPIRO, KB COX, S GLASSCO, JK JOHNSON, Sex Transm Dis 1992;19:341.

Single-dose cefixime versus single-dose ceftriaxone in the treatment of anti-microbial-resistant Neisseria gonorrhoeae infection

PJ PLOURDE, M TYNDALL, E AGOKI, et al J Infect Dis 1992;166:919.

Patterns of antibiotic susceptibility of gonococci isolated in Hong Kong, 1987-1990

km kam, cf lai, s egglestone, cbb chan, Sex Transm Dis 1992;19:284.

Comparative behavioural epidemiology of gonococcal and chlamydial infections among patients attending a Baltimore, Maryland, sexually transmitted disease clinic

EW HOOK, CA REICHART, DM UPCHURCH, P RAY, D CELENTANO, TC QUINN, Am J Epidemiol 1992:136:662.

Confidential HIV testing and condom promotion in Africa-impact on HIV and gonorrhoea rates

ALLEN, A SERUFILIRA, J BOGAERTS et al., JAMA 1992;268:3338.

Use of genomic fingerprinting in the characterization of Neisseria gonorrhoeae isolated in Valencia, Spain

M DASI, JM NOGUEIRA, JJ CAMARENA, et al. Eur J Clin Microbiol Infect Dis 1992;11:804.

Auxotyping of Neisseria gonorrhoeae as an additional epidemiological marker SK AGARWAL, M DEB, K PRAKASH, AK SHARMA, Ind 7 Med Res Sect A 1992;95:227.

Value of a DNA probe assay (Gen-probe) compared with that of culture for diagnosis of gonococcal infection

F VLASPOLDER, JAEM MUTSAERS, F BLOG, A NOTOWICZ, J Clin Microbiol 1992;31:107.

Evaluation of a fluorescent hybridization assay for the detection of Neisseria gonorrhoeae

RJ CANO, JC PALOMARES, MJ TORRES, RE KLEM, Eur J Clin Microbiol Infect Dis 1992;11:602.

Preliminary evaluation of the ligase chain reaction for specific detection of Neisseria gonorrhoeae

L birkenmeyer, as armstrong, J Clin Microbiol 1992;30:3089.

Effect of attachment factors (Pili Plus Opa) on Neisseria gonorrhoeae invasion of human fallopian tube tissue in vitroquantitation by computerized image

GL GORBY, N GB SCHAEFER, Microbial Pathogen 1992;13:93.

Binding of plasminogen to Neisseria meningitidis and Neisseria gonorrhoeae and formation of surface-associated plas-

M ULLBERG, P KEESELA, BE KRISTIANSEN, G KRON-VALL, J Infect Dis 1992;166:1329.

Antibodies to N-terminal peptides of gonococcal porin are bactericidal when gonococcal lipopolysaccharide is not sialylated

C ELKINS, NH CARBONETTI, VA VARELA, D STIRE-WAIT, DG KLAPPER, PF SPARLING, Mol Microbiol 1992:6:2617.

Regulation of catalase in Neisseria gonorrhoeae-effects of oxidant stress and exposure to human neutrophils

hy zheng, dj hassett, k bean, ms cohen, ${\mathcal J}$ ${\it Clin}$ Invest 1992;90:1000.

Effect of exogenous sialylation of the lipooligosaccharide of Neisseria gonorrhoeae on opsonophagocytosis

JJ KIM, DG ZHOU, RE MANDRELL, JM GRIFFISS, Infect Immun 1992;60:4439.

Identification and molecular analysis of a protein 63-kilodalton stress from Neisseria gonorrhoeae

y pannekoek, jpm vanputten, j dankert, ${\mathcal J}$ Bacteriol 1992;174:6928.

Isolation and nucleotide sequence of the gene (aniA) encoding the major anaerobically induced outer membrane protein of Neisseria gonorrhoeae

GT HOEHN, VL CLARK, *Infect* 1992;**60**:4695. Immun

The major anaerobically induced outer membrane protein of Neisseria gonorrhoeae. Pan-1, is a lipoprotein
GT HOEHN, VL CLARK, Infect

Immun 1992;60:4704.

Chlamydia

Chlamydial infection in Canada PR GULLY, Can Med Ass J 1992;147:893.

Association of genital infection with specific Chlamydia trachomatis serovars and race

KA WORKOWSKI, RF SUCHLAND, MB PETTINGER, WE STAMM., J Infect Dis 1992;166:1445.

Chlamydial genital infections and laparoscopic findings in infertile women

A LUCISANO, G MORANDOTTI, R MARANA, F LEONE, G BRANCA, S DELLACQUA, A SANNA, Eur J Epidemiol 1992;8:645.

Detection of Chlamydia trachomatis by the polymerase chain reaction in young patients with acute epididymitis

A ELEY, KM OXLEY, RC SPENCER, GR KINGHORN, ET BENAHMEIDA, CW POTTER, Eur J Clin Microbiol Infect Dis 1992;11:620.

Detection of cervical Chlamydia tra-chomatis and Neisseria gonorrhoeae with deoxyribonucleic acid probe assays in obstetric patients

IK HOSEIN, AM KAUNITZ, SJ CRAFT, Am J Obstet Gynecol 1992;167:588.

Evaluation of new anti-infective drugs for the treatment of sexually transmitted chlamydial infections and related clinical syndromes

HH HANDSFIELD, AR RONALD, L COREY, JA MCCUTCHAN, Clin Infect Dis 1992;15:S131.

Compliance with antibiotic therapy for Chlamydia trachomatis and Neisseria gonorrhoeae

BP KATZ, BW ZWICKI, VA CAINE, RB JONES, Sex Transm Dis 1992;19:351.

Efficacy of single-dose azithromycin versus doxycycline in the treatment of cervical infections caused by Chlamydia trachomatis

JM OSSEWAARDE, FHF PLANTEMA, M RIEFFE, RP NAWROCKI, A DEVRIES, AM VANLOON, Eur J Clin Microbiol Infect Dis 1992;11:693.

Azithromycin-induced block of elementary body formation in Chlamydia trachomatis

Antimicrob Agents ENGEL, 1992;36:2304.

Underdiagnosis of Chlamydia trachomatis infection—diagnostic limitations in patients with low-level infection

JSL LIN, WE JONES, LL YAN, et al. Sex Transm Dis 1992;19:259.

Nonculture tests for genital tract Chlamydia infection—what does the package insert mean and will it mean the same thing tomorrow

J SCHACHTER, WE STAMM, MA CHERNESKY, et al, Sex Transm Dis 1992;19:243.

Multicenter evaluation of the antigen Chlamydia enzyme immunoassay for diagnosis of Chlamydia trachomatis genital infection

A CLARK, WE STAMM, C GAYDOS, et al., J Clin Microbiol 1992;30:2762.

Comparison of two culture approaches, blind passage and dual observation, for detecting Chlamydia trachomatis in various prevalence populations

SJ ZIMMERMAN, E MOSES, N SOFAT. WR BARTHOLOMEW, D AMSTERDAM., \mathcal{J} Clin Microbiol 1992:30:2938.

Comparison of cytobrushes with swabs for recovery of endocervical cells and for Chlamydiazyme detection of Chlamydia trachomatis

JA KELLOGG, JW SEIPLE, JL KLINEDINST, JS LEVISKY, 7 Clin Microbiol 1992;30:2988.

Effects of broadening the gold standard on the performance of a chemiluminometric immunoassay to detect Chlamydia trachomatis antigens in centrifuged first void urine and urethral swab samples from men

D JANG, IW SELLORS, IB MAHONY, L PICKARD, MA CHERNESKY, Sex Transm Dis 1992;19:315.

Evaluation of a chemiluminometric immunoassay for detection of *Chlamydia trachomatis* in the urine of male and female patients

C SCIEUX, A BIANCHI, S HENRY, et al., Eur J Clin Microbiol Infect Dis 1992;11:704.

Detection of *Chlamydia trachomatis* in endocervical specimens by polymerase chain reaction

MJ LOEFFELHOLZ, CA LEWINSKI, SR SILVER, et al. J Clin Microbiol 1992;30:2847.

Establishment of a particle-counting method for purified elementary bodies of chlamydiae and evaluation of sensitivities of the IDEIA Chlamydia kit and DNA probe by using the purified elementary bodies

N MIYASHITA, A MATSUMOTO, J Clin Microbiol 1992;30:2911.

Identification of Chlamydia trachomatis antigens by use of murine T-cell lines
PR BEATTY, RS STEPHENS, Infect Immun

1992;**60**:4598.

Gene typing of *Chlamydia trachomatis* by polymerase chain reaction and restriction endonuclease digestion

CA GAYDOS, L BOBO, L WELSH, EW HOOK, R VISCIDI, TC QUINN, Sex Transm Dis 1992;19:303.

Inhibition of *Chlamydia trachomatis* growth in mouse fibroblasts by liposome-encapsulated tetracycline

H ALAWADHI, GV STOKES, M REICH, J Antimicrob Chemother 1992;30:303.

Effects of ascorbic acid on *Chlamydia trachomatis* infection and on erythromycin treatment in primary cultures of human amniotic cells

SK WANG, DL PATTON, CC KUO, J Clin Microbiol 1992;30:2551.

Effect of gamma-interferon on resolution of murine chlamydial genital infection

RG RANK, KH RAMSEY, EA PACK, DM WILLIAMS, Infect Immun 1992;60:4427.

Evaluation of the humoral immune response in trachoma to *Chlamydia trachomatis* major outer membrane proteins by sequence-defined immunoassay

HM JONES, J SCHACHTER, RS STEPHENS, J Infect Dis 1992;166:915.

Pelvic Inflammatory Disease

A longitudinal study of pelvic inflammatory disease

CM STACEY, PE MUNDAY, D TAYLOR ROBINSON, Br J Obstet Gynaecol 1992;99:994.

The intrauterine device, pelvic inflammatory disease and infertility—the confusion between hypothesis and knowledge DA GRIMES, Fert Steril 1992;58:670.

Current cigarette smoking and risk of acute pelvic inflammatory disease

D SCHOLES, JR DALING, AS STERGACHIS, Am J Public Health 1992;82:1352.

Evaluation of new anti-infective drugs for the treatment of acute pelvic inflammatory disease

RL SWEET, JG BARTLETT, DL HEMSELL, JS SOLOMKIN, F TALLY, Clin Infect Dis 1992;15:853.

Microbial etiology of urban emergency department acute salpingitis—treatment with ofloxacin

DE SOPER, NJ BROCKWELL, HP DALTON, Am J Obstet Gynecol 1992;167:653.

Candidiasis

Histopathology of human vaginal candiosis

VL BYKOV, Mycoses 1992;35:77.

Genetic similarity of Candida albicans strains from vaginitis patients and their partners

J SCHMID, M ROTMAN, B REED, CL PIERSON, DR SOLL, J Clin Microbiol 1992;31:39.

Day-to-day follow-up after a short oral treatment of acute vaginal candidosis with itraconazole

GPJ BEYER, HJ VOORHOEVEDENHARTOG, Mycoses 1992;35:99.

Azole drug resistance in Candida species DW WARNOCK, J Med Microbiol 1992;37:225.

Influence of alkaline pH on the direct lethal action of miconazole against Candida albicans

WH BEGGS, Mycopathol 1992;120:11.

Modulation of interactions of *Candida* albicans and endothelial cells by fluconazole and amphotericin

MA GHANNOUM, SG FILLER, AS IBRAHIM, Y FU, JE EDWARDS, Antimicrob Agents Chemother 1992;36:2239.

Identification of a mannoprotein fraction from Candida albicans that enhances human polymorphonuclear leukocyte (PMNL) functions and stimulates lactoferrin in PMNL inhibition of candidal growth

C PALMA, D SERBOUSEK, A TOROSANTUCCI, A CASSONE, JY DJEU, *J Infect Dis* 1992;**166**:1103.

Comparison of molecular typing methods for Candida albicans

PT MAGEE, L BOWDIN, J STAUNDINGER, J Clin Microbiol 1992;30:2674.

Yeast-specific DNA probes and their application for the detection of Candida albicans

ar holmes, yc lee, rd cannon, hf jenkinson, mg shepherd, \emph{J} Med Microbiol 1992; $\emph{37}$:346.

Typing of *Candida albicans* **strains**FC ODDS, DL BRAWNER, J STAUDINGER, PT MAGEE,

DR SOLL, J Med Vet Mycol 1992;30:87.

Molecular basis of Candida albicans

adhesions
MJ KENNEDY, RA CALDERONE, JE CUTTER, et al.,
J Med Vet Mycol 1992;30:S1:95.

Bacterial vaginosis

Frequency of mobiluncus spp in bacterial vaginosis in Italy

M FENOCCHI, M GATTI, Microbiologica 1992;15: 409.

Trichomoniasis

Risk assessment and laboratory diagnosis of trichomoniasis in men

JN KRIEGER, M VERDON, N SIEGAL, C CRITCHLOW, KK HOLMES, *J Infect Dis* 1992;166:1362.

Successful response of metronidazoleresistant trichomonal vaginitis to tinidazole—a case report

KA HAMED, AE STUDEMEISTER, Sex Transm Dis 1992;19:339.

Split-dose metronidazole or single-dose tinidazole for the treatment of vaginal trichomoniasis

P OPRASERTSAWAT, T JETSAWANGSRI, Sex Transm Dis 1992;19:295.

A novel neutrophil-activating factor released by *Trichomonas vaginalis*MF SHAIO, PR LIN, CS LEE, SC HOU, P TANG, KD

YANG, Infect Immun 1992;60:4475.

Syphilis and other treponematoses

Is syphilis an incurable disease

RC GHINNSBERG, Y NITZAN, Med Hypotheses 1992;39:35.

Osseous syphilis

P DISDIER, JR HARLE, L SWIADER, D GAMBARELLI, G MOULIN, PJ WEILLER, Arthritis Rheum 1992;35:1241.

Evaluation of new anti-infective drugs for the treatment of syphilis

AR RONALD, M SILVERMAN, JA MCCUTCHAN, L COREY, HH HANDSFIELD, Clin Infect Dis 1992;15:S140.

Management of syphilis in human immunodeficiency virus-infected patients EW HOOK, Am J Med 1992;93:477.

Response of latent syphilis or neurosyphilis to ceftriaxone therapy in persons infected with human immunodeficiency virus

ME DOWEKK, PG ROSS, DM MUSHER, TR CATE, RE BAUGHN, $Am\ \mathcal{J}\ Med\ 1992; 93:481$.

High-risk behaviours for transmission of syphilis and human immunodeficiency virus among crack cocaine-using women—a case study from the Midwest HA SIEGAL, RG CARLSON, R FALCK, MA FORNEY, JC WANG, LLI., Sex Transm Dis 1992;19:266.

Behaviours of crack cocaine users and their impact on early syphilis intervention J GREENBERG, D SCHNELL, R CONLON, Sex Transm Dis 1992;19:346.

Characteristics of mothers of live infants with congenital syphilis in Florida, 1987-1989

JCA DESENCLOS, M SCAGGS, JE WROTEN Am J Epidemiol 1992;136:657.

Treponemal infection and the outcome of pregnancy in a rural area of the Gambia, West Africa

AM GREENWOOD, V DALESSANDRO, F SISAY, BM GREENWOOD, J Infect Dis 1992;166:842.

IgM antibody to Treponema pallidum in cerebrospinal fluid of infants with congenital syphilis

PJ SANCHEZ, GD WENDEL, MV NORGARD, Am J Dis Child 1992;146:1171.

Congenital syphilis in the past—slaves at Newton plantation, Barbados, West Indies KP JACOBI, DC COOK, RS CORRUCCINI, JS HANDLER, Am J Physical Anthropol 1992;89:145.

Comparison of the fluorescent treponemal antibody absorption (FTA-ABS) test with the FTA-ABS double staining test for detection of antitreponemal immunoglobulin-M in the 19S fraction of human serum

G OZANNE, MA DHALEWYN, SA LARSEN, J Clin Microbiol 1992;31:102.

Herpes

The management of pregnancies complicated by genital infections with herpes simplex virus

CG PROBER, L COREY, ZA BROWN, et al., Clin Infect Dis 1992;15:1031.

The placental transport and use of acyclovir in pregnancy

CV FLETCHER, J Lab Clin Med 1992;120:821.

Acyclovir transport by the human placen-

GI HENDERSON, ZQ HU, RF JOHNSON, AB PEREZ, YQ YANG, S SCHENKER, J Lab Clin Med 1992;120:885.

Effect of acyclovir on selected immune functions

M KANKOVAVANCUROVA, M JILEK, V FISTERIE, J PROCHAZKOVA, Acta Virol 1992;36:284.

Congenital neonatal herpes simplex virus infection

B SARKELL, WK BAYLOCK, H VERNON, $\it J$ Am Acad Dermatol 1992;27:817.

Underdiagnosis of genital herpes J KAPLAN, N Engl J Med 1992;327:1098.

A 15-year surveillance study of antibodies to herpes simplex virus type-1 and type-2 in a cohort of young girls

B CHRISTENSON, M BOTTIGER, A SVENSSON, S JEANSSON, J Infect 1992;25:147.

Prevalence and correlates of herpes simplex infections—the population-based AIDS in multiethnic neighborhoods study D SIEGEL, E GOLDEN, AE WASHINGTON, et al., JAMA 1992;268:1702.

Evaluation of new anti-infective drugs for the treatment of genital infections due to herpes simplex virus

L COREY, JA MCCUTCHAN, AR RONALD, HH HANDS-FIELD, Clin Infect Dis 1992;15:S99.

Mucosal delivery of herpes simplex virus vaccine

JC BOWEN, HO ALPAR, R PHILLPOTTS, MRW BROWN, Res Virol 1992;143:269.

Rapid DNA diagnosis of herpes simplex virus serotypes

T MATSUMOTO, O YAMADA, A ITAGAKI, S ISHIDA, T KAMAHORA, T KURIMURA, J Virol Meth 1992;40:119.

The latency-associated transcripts of herpes simplex virus—RNA in search of function

NW FRASER, TM BLOCK, JG SPIVACK, *Virol* 1992;191:1.

Herpes simplex type-1 and type-2—latency in the genital tract of guinea pigs ML LANDRY, C BULL, Intervirol 1992;33:204.

Herpesvirus serology, aberrant specific immunoglobulin-G2 and immunoglobulin-G3 subclass patterns and Gm allotypes in individuals with low levels of IgG3

A LINDE, R SODERSTROM, CIE SMITH, et al., Clin Exp Immunol 1992;90:199.

Disulfide bond structure of glycoprotein-D of herpes simplex virus type-1 and type-

D LONG, WC WILCOX, WR ABRAMS, GH COHE, RJ EISENBERG, J Virol 1992;66:6668.

Herpes simplex virus-infected human fibroblasts are resistant to and inhibit cytotoxic lymphocyte-T activity

CM POSAVAD, KL ROSENTHAL, J Virol 1992;66:6264.

Human papillomavirus infection

Phylogenetic classification of human papillomaviruses—correlation with clinical manifestations

M VANRANST, JB KAPLAN, RD BURK, J Gen Virol 1992;73:2653.

The causal link between human papillomavirus and invasive cervical cancer—a population-based case-control study in Colombia and Spain

N MUNOZ, FX BOSCH, S DESANJOSE, et al. Int J Cancer 1992;52:743.

Risk factors for cervical cancer in Colombia and Spain

FX BOSCH, N MUNOZ, S DESANJOSE, et al. Int J Cancer 1992;52:750.

Risk factors inducing the persistence of high-risk genital papillomaviruses in the normal cervix

C VANDENVELDE, D VANBEERS, J Med Virol 1992;38:226.

Oral contraceptive use and human papillomavirus infection in women without abnormal cytological results

T CSIKYMASZAROS, J CZEGLEDY, L GERGELY., Med Microbiol Immunol 1992;181:181.

The relationship between contraceptives, sexual practices and cervical human papillomavirus infection among a college population

BJ BURKETT, CM PETERSON, LM BIRCH, et al. J Clin Epidemiol 1992;45:1295.

Anal intraepithelial neoplasia—part of a multifocal disease process

JH SCHOLEFIELD, WGE HICKSON, JHF SMITH, K ROGERS, F SHARP, *Lancet* 1992;**340:**1271.

Natural history of anal cytologic abnormalities and papillomavirus infection among homosexual men with group-IV

JM PALEFSKY, EA HOLLY, J GONZALES, K LAMBORN, H HOLLANDER, \mathcal{J} Acq Immun Def Syn 1992;5:1258.

Viral co-infections in human papillomavirus-associated anogenital lesions according to the serostatus for the human immunodeficiency virus

C BERNARD, C MOUGIN, L MADOZ, et al. Int J Cancer 1992;52:731.

Genotypes and sequence variants of human papillomavirus DNAs from human immunodeficiency virus type-I-infected women with cervical intraepithelial neoplasia

JP ICENOGLE, M LAGA, D MILLER, AT MANOKA, RA TUCKER, WC REEVES, J Infect Dis 1992;166:1210.

A cohort study of the risk of cervical intraepithelial neoplasia grade-2 or grade-3 in relation to papillomavirus infection LA KOUTSKY, KK HOLMES, CW CRITCHLOW, et al., N Engl J Med 1992;327:1272.

Lymphocyte subset and NK activity in cervical intra-epithelial neoplasia

GG GARZETTI, M CIGNITTI, A CIAVATTINI, N FABRIS, C ROMANINI., Gynecol Obstet Invest 1992;34:119.

Adjuvant treatment of recalcitrant genitoanal warts with systemic recombinant interferon-alpha-2c

RM HOPFL, M SANDBICHLER, BWH ZELGER, FG CONRAD, PO FRITSCH, Acta Derm Vener 1992;72:383.

Southern blot hybridization and PCR in detection of oral human papillomavirus (HPV) infections in women with genital HPV infections

JK KELLOKOSKI, SM SYRJANEN, F CHANG, M YLISKOSKI, KJ SYRJANEN *J Oral Pathol Med* 1992;21:459.

Detection of human papillomavirus type-16 DNA in oral squames from normal young adults

H JALAL, CM SANDERS, SS PRIME, C SCULLY, NJ MAITLAND, J Oral Pathol Med 1992;21:465.

Human papillomavirus type-6 infection involving cutaneous nongenital sites

A BLAUVELT, AM DUARTE, C PRUKSACHATKU-NAKORN, CL LEONARDI, LA SCHACHNER, J Am Acad Dermatol 1992;27:876.

Human papillomavirus genomes in male urethral cells

G DELLATORRE, R DONGHI, POD LIMA, et al. Am \mathcal{J} Pathol 1992;141:1181.

High frequency of human papillomavirus infection in carcinoma of the urinary bladder

K ANWAR, H NAIKI, K NAKAKUKI, M INUZUKA, Cancer 1992;70:1967.

Presence and physical state of HPV DNA in prostate and urinary-tract tissues

A ROTOLA, P MONINI, D DILUCA, et al. Int J Cancer 1992;52:359.

Detection of human papillomavirus in the prostate by polymerase chain reaction and in situ hybridization

GK IBRAHIM, PE GRAVITT, KL DITTRICH, et al. J Urol 1992;148:1822.

The predictive value of human papillomavirus (HPV) typing in the prognosis of bronchial squamous cell papillomas.

HH POPPER, G WIRNSBERGER, FM JUTTNERSMOLLE, MG PONGRATZ, M SOMMERSGUTTER, *Histopathol* 1992;21:323.

Colposcopic and histologic findings and human papillomavirus (HPV) DNA test variability in young women positive for HPV DNA

AB MUSCICKI, JM PALEFSKY, J GONZALES, G SMITH, GK SCHOOLNIK, J Infect Dis 1993;166:951.

Predictive value of human papillomavirus DNA detection by filter hybridization and polymerase chain reaction in women with negative results of colposcopic examination

GJ NUOVO, J MORITZ LL WALSH, P MACCONNELL, J KOULOS, Am J Clin Pathol 1992;**98:4**89.

Polymerase chain reaction detection of human papillomavirus—quantitation may improve clinical utility

EAB MORRISON, GL GOLDBERG, AS KADISH, RD BURK, J Clin Microbiol 1992;30:2539.

Comparison of ViraPap, Southern hybridization and polymerase chain reaction methods for human papillomavirus identification in an epidemiological investigation of cervical cancer

E GUERRERO, RW DANIEL, FX BOSCH, et al, J Clin Microbiol 1993;30:295.

Comparative analysis of human papillomavirus detection by dot blot hybridization and non-isotopic in situ hybridization G TRONCONE, SM ANDERSON, CS HERRINGTON, et al. J Clin Pathol 1992;45:866.

In situ hybridization for the detection of human papillomavirus (HPV) in gynaecological biopsies—a study of 2 commercial kits

I ZEHBE, E RYLANDER, A STRAND, E WILANDER, Anticancer Res 1992;12:1383.

Biologic characteristics of specific human papillomavirus types predicted from morphology of cervical lesions

AS KADISH, RJ HAGAN, DB RITTER, et al, Hum Pathol 1992;23:1262.

Alterations of the p53 gene in human primary cervical carcinoma with and without human papillomavirus infection

M FUJITA, M INOUE, O TANIZAWA, S IWAMOTO, T ENOMOTO, Cancer Res 1992;52:5323.

Expression of the L2 and E7 genes of the human papillomavirus type-16 in female genital dysplasia

E AUVINEN, H KUJARI, P ARSTILA, V HUKKANEN, Am J Pathol 1992;141:1217.

Detection of capsid antigen of human papillomavirus (HPV) in benign lesions of female genital tract using anti-HPV monocloncal antibody

T IWASAKI, T SATA, M SUGASE, et al. J Pathol 1992;168:293.

Enhanced tenascin expression in cervical and vulvar koilocytotic lesions

O TIITTA, I WAHLSTROM, J PAAVONEN, A LINNALA, S SHARMA, VE GOULD, I VIRTANEN, Am J Pathol 1992;141:907.

Isolation of 2 keratinocyte cell lines derived from HPV-positive dysplastic vaginal lesions

S HIETANEN, E AUVINEN, S GRENMAN, et al, Int J Cancer 1992;52:391.

Papillomavirus LI major capsid protein self-assembly into virus-like particles that are highly immunogenic

R KIRNBAUER, F BODY, N CHENG, DR LOWY, JT SCHILLER, *Proc Nat Acad Sci USA* 1992;**89**:12180.

Progression from papilloma to carcinoma is accompanied by changes in antibody response to papillomavirus proteins

YL LIN, LA BORENSTEIN, R SELVAKUMAR, R AHMED, FO WETTSTEIN, J Virol 1992;67:382.

Cellular transformation by a unique isolate of human papillomavirus type-11

RC MCGLENNEN, J GHAI, RS OSTROW, K LABRESH, JF SCHNEIDER, AJ FARAS, *Cancer Res* 1993;52:5872.

Presence of ras oncogene mutations and human papillomavirus DNA in human prostate carcinomas

K ANWAR, K NAKAKUKI, T SHIRAISHI, H NAIKI, R YATANI, M INUZUKA, Cancer Res 1993;52:5991.

Sequential combined tumorigenic effect of HPV-16 and chemical carcinogens

SL LI, MS KIM, HM CHERRICK, J DONIGER, NH PARK, Carcinogenesis 1992;13:1981.

Tamoxifen stimulates human papillomavirus type-16 gene expression and cell proliferation in a cervical cancer cell line JY HWANG, BY LIN, FM TANG, WCY YU, Cancer Res 1992;52:6848.

Preferential chromosome loss in human papillomavirus DNA-immortalized mammary epithelial cells

K SWISSHELM, M LEONARD, R SAGER, Gene Chromosome Cancer 1993;5:219.

Down-regulation of keratin-14 gene expression after v-Ha-ras transfection of human papillomavirus-immortalized human cervical epithelial cells

PE BOWDEN, CD WOODWORTH, J DONIGER, JA DIPAOLO, Cancer Res 1993;52:5865.

Autocrine growth limitation of human papillomavirus-type-16 harboring keratinocytes by constitutively released tumor necrosis factor-α

J MALEJCZYK, M MALEJCZYK, A KOCK, et al. J Immunol 1992;149:2702.

Human papillomavirus type-16 expresses a variety of alternatively spliced messenger RNAs putatively encoding the E2 protein

L SHERMAN, N ALLOUL., Virology 1992;191:953.

Cervical cytology and colposcopy

Mild cervical cytological abnormalities MI SHAFI, DM LUESLEY, JA JORDAN, BMJ 1992;305:1040.

Oral contraceptive use and cervical intraepithelial neoplasia

AL COKER, MF MCCANN, BS HULKA, LA WALTON, J Clin Epidemiol 1992;45:1111.

The Bethesda system—a proposal for reporting abnormal cervical smears based on the reproducibility of cytopathologic diagnoses

ME SHERMAN, MH SCHIFFMAN, YS EROZAN, S WACHOLDER, RJ KURMAN, Arch Pathol Lab Med 1992;116:1155.

Efficacy of lignocaine analgesia during treatment of the cervix

KE ROGSTAD, DJ WHITE, IH AHMEDJUSHUF, Lancet 1992;340:942.

Invasive cervical cancer following cryotherapy for cervical intraepithelial neoplasia or human papillomavirus infection.

C SCHMIDT, RG PRETORIUS, M BONIN, L HANSON, N SEMRAD, W WAIRING, *Obstet Gynecol* 1992; **80**:797.

Histological incomplete excision of CIN after large loop excision of the transformation zone (LLETZ) merits careful follow-up, not retreatment

JB MURDOCH, PR MORGAN, A LOPES, JM MONAGHAN, $Br\ \mathcal{F}$ Obstet Gynaecol 1992;99:990.

3-group metaphase as a morphologic criterion of progressive cervical intraepithelial neoplasia

MJE MOURITS, WJLM PIETERS, H HOLLEMA, MPM BURGER, Am J Obstet Gynecol 1992;167:591.

Oestrogen receptor message in premalignant and normal cervical cells—a methodological study

HC KITCHENER, L NEILSON, JCM MACNAB, Eur J Cancer 1992;29A:252.

Other sexually transmitted diseases

Heterosexual transmission of hepatitis-C virus in family groups without risk factors GM PEANO, LM FENOGLIO, G MENARDI, R BALBO, D MARENCHINO, S FENOGLIO, BMJ 1992;305:1473.

Infection by hepatitis-B and hepatitis-C virus in nonintravenous drug using female prostitutes in Spain

A ORDUNA, MA BRATOS, P GUTIERREZ, et al., Eur J Epidemiol 1992;8:656.

Seroepidemiologic study of hepatitis-C virus in sexually transmitted disease risk groups.

A STARY, W KUPP, H HOFMANN, C HELLERVITOUCH, C KUNZ, Sex Transm Dis 1992;19:252.

Sexual transmission of hepatitis-C virus among female prostitutes and patients with sexually transmitted diseases in Fukuoka, Kyushu, Japan

K NAKASHIMA, S KASHIWAGI, J HAYASKI, et al., Am J Epidemiol 1992;136:1132.

Lack of protective immunity against reinfection with hepatitis-C virus

P FARCI, HJ ALTER, S GOVINDARAJAN, et al. Science 1992;258:135.

Hepatitis-C and non-A-hepatitis non-B-hepatitis revisited—hepatitis-E, hepatitis-F and hepatitis-G

J CRASKE, J Infect 1992;25:243.

Treatment of chancroid with a single dose of spectinomycin

M GUZMAN, J GUZMAN, M BERNAL, Sex Transm Dis 1992;19:291.

Evaluation of new anti-infective drugs for the treatment of chancroid

AR RONALD, L COREY, JA MCCUTCHAN, HH HANDS-FIELD, Clin Infect Dis 1992;15:S108.

Association of *Haemophilus ducreyi* with cell-culture lines

L SHAH, HA DAVIES, RA WALL, J Med Microbiol 1992;37:268.

Use of an absorption enzyme immunoassay to evaluate the *Haemophilus ducreyi* specific and cross-reactive humoral immune response to humans

MJ ALFA, N OLSON, P DEGAGNE, et al. Sex Transm Dis 1992;19:309.

Public health and social aspects

Commentary: methods women can use that may prevent sexually transmitted disease, including HIV

MJ ROSENBERG, EL GOLLUB, Am J Public Health 1992;82:1473.

Multiple partners and partner choice as risk factors for sexually transmitted disease among female college students

GP JOFFE, B FOXMAN, AJ SCHMIDT, et al. Sex Transm Dis 1992;19:272.

Prior episode of sexually transmitted disease and subsequent sexual risk-reduction practices—a need for improved risk-reduction interventions

P OCAMPO, M DEBOER, RR FADEN, N KASS, AC GIE-LEN, J ANDERSON, Sex Transm Dis 1992;19:326.

Condoms for men, not women-results of brief promotion programs

DA COHEN, C DENT, D MACKINNON, G HAHN, Sex Transm Dis 1992;19:245.

Contraception and prevalence of sexually transmitted diseases among adolescents and young adults in Uganda

WKA AGYEI, EJ EPEMA, M LUBEGA, Int 7 Epidemiol 1992;21:981.

Condom use and sexual habits of heterosexual intravenous drug users in Northern

D SERRAINO, S FRANCESCHI, E VACCHER, et al, Eur 7 Epidemiol, 1992;21:981.

Microbiology and Immunology

Local immunity in the male reproductive

VL CLIFTON, AJ HUSBAND, DJ KAY, Immun Cell Biol 1992;70:301.

Density and distribution of Langerhans cells in the human uterine cervix

AE MORRELLI, G DIPAOLA, L FAINBOIM, Arch Gynecol Obstet 1992;252:65.

The balance of macrophage subsets may be customised at mucosal surfaces

C HUTTER, LW POULTER, FEMS Microbiol Immunol 1992;105:309.

Comparison of two antigen assays for rapid intrapartum detection of vaginal group-B streptococal colonisation

M GREEN, B DASHEFSKY, ER WALD, S LAIFER, J HARGER, R GUTHRIE, J Clin Microbiol 1992;31:78

Dermatology

Fusidic acid cream in the treatment of plasma cell balanitis

CS PETERSEN, K THOMSEN, J Am Acad Dermatol 1992;27:633.

Crohn's disease of the vulva

NI KIM, JY EOM, WY SIM, CR HAW, J Am Acad Dermatol 1992;27:764.

Crohn's disease presenting as unilateral labial hypertrophy

SL WERLIN, NB ESTERLY, H OECHLER, J Am Acad Dermatol 1992;27:893.

Latex allergy-what do we know JE SLATER, J Allerg Clin Immunol 1992;90:279.

Neurofibromatosis of the vulva

FM LEWIS, MS LEWIS JONES, PG TOON, IP ROLLASON. Br $\mathfrak F$ Dermatol 1992;127:540.

Steroid-responsive pyoderma gangreno-sum with vulvar and pulmonary involvement.

C LEBBE, I MOULONGUETMICHAU, PPERRIN, F BLANC, J FRIJA, J CIVATTE, J Am Acad Dermatol 1992;27:623.

Miscellaneous

Special issues in clinical trials of new antiinfective drugs for the treatment of sexually transmitted diseases

HH HANDSFIELD, JA MCCUTCHAN, L COREY, AR RONALD, Clin Infect Dis 1992;15:S96.

Evaluation of new anti-infective drugs for the treatment of vaginal infections

JA MCCUTCHAN, AR RONALD, L COREY, HH HANDS-FIELD, Clin Infect Dis 1992;15:S115.

The urethral syndrome—why what we do works or doesn't

GM TESTA, Med J Australia 1992;157:549.

Management of chronic urethral symptoms in men

MH AUGENBRAUN, M CUMMINGS, WM MCCOR-MACK, Clin Infect Dis 1992;15:714.

The characterization of nonbacterial prostatitis-search for an etiology.

LMD SHORTLIFFE, RG SELLERS, J SCHACHTER, ${\mathcal J}$ Urol 1992:148:1461.

Psychosexual aspects of the evaluation and management of vulvar vestibulitis LR SCHOVER, DD YOUNGS, R CANNATA, Am J Obstet Gynecol 1992;167:630.

Results of surgical management of Peyronie's disease

PD ODONNELL, J Urol 1992;148:1184.

of colorectal diseasestransmitted diseases and anal papillomas. DJ JONES, BP GOORNEY, BMJ 1992;305:820.

Prevalence of sexually transmitted disease infection in women alleging rape AG DAVIES, JC CLAY, Sex Transm Dis 1992;19:298.

Sexual assault-review of a national model protocol for forensic and medical evaluation

WW YOUNG. AC BRACKEN, MA GODDARD, S MATHESON, Obstet Gynecol 1992;80:878.

Chronically evolving viral hepatitis Arch Virol 1992 Suppl 4 (whole issue)

Adverse events after hepatitis-B vaccina-

P DUCLOS, Can Med Assoc 3 1992;147:1023.

Revised guidelines for booster vaccination against hepatitis-B

D HOLTON, Can Med Assoc J 1992;147:1029.

Sexual behaviour in Britain and France J BANCROFT

BMJ 1992;305:1447.

Sebaceous glands and hair follicles in the cervix uteri

MC ROBLEDO, JJ VAZQUEZ, F CONTRERASMEJUTO, G LOPEZGARCIA, Histopathology 1992;21:278.